



Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil

Francisco I. de Souza¹, Nildo da S. Dias², Osvaldo N. de Sousa Neto¹, Ana Claudia Medeiros de Souza¹, Raniere B. de Lira¹ and Francisco Souto de Souza Junior¹

¹F Federal University of Semi-Arid, Mossoró, Brazil.

²Federal University of Semi-Arid. BR 110 km 47. 59625-900. Mossoró, RN, Brazil.
nildo@ufersa.edu.br. Corresponding author

ABSTRACT

The aim of this research was to study the feasibility of growing organic vegetables using saline waste from the water treatment station in the community Bom Jesus, in Campo Grande, RN. The research began with the establishment of a community garden according to systematic principles of agricultural sustainability in ecological basis. The participatory planning, in which all activities in the project were planned and performed democratically with the community, was the methodology adopted. Vegetables like coriander, chive, tomato and rocket were grown in the site available for the research, sowing a legumes and grasses mixture to produce plant material, a breeding (elephantgrass), and planting forest and halophyte species (*Atriplex nummularia*). The appropriate disposal of the saline waste gave a conservative nature to the project and also the possibility of an alternative source of water for agriculture, emphasizing the efficiency of use of water in semiarid regions. It is concluded that there were yield losses in all species grown with water of saline waste – from 25.30% to 26.7% in relation to the conventional irrigation systems (fresh water); but there was biomass. However, the vegetable growth was successful in the community, because it could change the resident's mind about the disposal and reuse of the saline waste.

Keywords: Action research, Saline waste, Water resources, Green manure, Brazil.

1. INTRODUCTION

The families in the community Bom Jesus have historically fought and resisted against natural climatic factors. The limited and poor water resources is a problem faced by the community, where there are only four small reservoirs to provide water for domestic use and for the animals until a certain period of years and the residents suffer from water shortage during the dry months. To solve this problem, the Brazilian government

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ “Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil”. EFITA-WCCA-CIGR Conference “Sustainable Agriculture through ICT Innovation”, Turin, Italy, 24-27 June 2013. The authors are solely responsible for the content of this technical presentation. The technical presentation does not necessarily reflect the official position of the International Commission of Agricultural and Biosystems Engineering (CIGR) and of the EFITA association, and its printing and distribution does not constitute an endorsement of views which may be expressed. Technical presentations are not subject to the formal peer review process by CIGR editorial committees; therefore, they are not to be presented as refereed publications.

Sustainable Agriculture through ICT innovation

conducted the program “Água Boa” according to which a water treatment station by reverse osmosis was installed in the community as alternative for the families to have drinking water (Mohamed et al, 2005, Dias et al., 2011).

Historically, due to its high cost, desalination was limited to applications as a source for potable water and for high-quality water for industrial uses. However, technological advances in the last years have driven down the cost of desalinated water due to reductions in the price of equipment, reduction in energy demand and know-how on water treatment (Tsiourtis, 2001, Fritzmann et al., 2007). In Rio Grande do Norte State, the first desalination plant experience began in 1997 and since then, the numbers of desalination plants have been increased annually.

The community has been working to build a pipe for water supply in order to solve the problem of water shortage. Meanwhile, the desalination and the tanks are the alternative to ensure domestic water supply. However, the residues (concentrate and brine) resulting from the water desalination do not pass by any treatment or appropriate disposal. They are disposed directly on soil and into the water, causing problems of salinity in soil and affecting water quality in dams and rivers next to the community (Tchouaffe, 2007) it is important to know how salt contamination affects crops. Water is taken up by the fine roots of the crops through osmosis, which involves the movement of water from the area of low salt concentration (such as the soil) to area of high salt concentration (such as inside root cells). When salt concentrations are high, the movement of water from the soil to the root is slowed down. When the salt concentrations in the soil are higher than inside the root cells, the soil will draw water from the root, and the root, and the crop will wilt and sometimes die. This is the basic way in which salinization affects crop production.

Generally brine does not receive any treatment in Brazil; anyway, it has been poured on soil, providing high accumulation of salts in soils (Porto et al., 2001) causing short term salinization problems in communities that benefit from the desalination technology.

The families concerned about the environmental problems caused by the disposal of saline waste resulting from the water treatment tried some alternatives, but all with very high costs. The Federal University of Semi-arid and the technical assistance of the Nucleo Sertão Verde, in partnership with Program of Undergraduate Research and Technology for Micro and Small Enterprises (*Programa de Iniciação Científica e Tecnológica para Micro e Pequenas Empresas – BITEC 2009*), presented a proposal for a better destiny of wastewater from desalination.

Thus, the aim of this action research was to study the feasibility of using saline waste from the water treatment to support the production of organic vegetables in the community Bom Jesus, Campo Grande, RN. The research began with the establishment

P0074

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ “Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil”. EFITA-WCCA-CIGR Conference “Sustainable Agriculture through ICT Innovation”, Turin, Italy, 24-27 June 2013.

Sustainable Agriculture through ICT innovation

of a community garden according to systematic principles of agricultural sustainability in ecological basis.

2. MATERIAL AND METHODS

This study was conducted from July to December 2009 with the establishment of a community garden according to systematic principles of agricultural sustainability in ecological basis. The participatory planning, in which all activities in the project were planned and performed democratically with the community, was the methodology adopted. The group already had the basics of vegetable production, but needed specific training. As it is an innovative project, technicians from UFERSA and NUCLEO SERTA0 VERDE guided them.

Based on the site analysis, it was chosen an area measuring 12 x 20 m next to the desalination (Figure 1), where it was necessary to build a barbed wire fence – to isolate this area which is central to the community, a nursery for planting seedlings in beds, and a small irrigation system using waste water from water desalination processes to support the desalination of crops.



Figure 1. Salt water treatment station and disposal of saline waste in soil and water course

P0074

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ “Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil”. EFITA-WCCA-CIGR Conference “Sustainable Agriculture through ICT Innovation”, Turin, Italy, 24-27 June 2013.

Sustainable Agriculture through ICT innovation

Firstly, there was the site preparation for planting, and the construction of ‘smart beds’ with alternative system for water distribution on subsurface (Figure 2A) in order to avoid the contact of leafy vegetables with the saline waste. The beds were filled with a great amount of plant material (coconut fiber, plant debris, wood remains etc.), once the principle of reduction of matrix soil potential was tried to decrease the deleterious effects of saline waste water on plants. Besides, a legume and grass seed mixture was used in half of the available site (Table 1) for the production of biomass which later served as green manure and organic material in soil.

Table 1. Composition and quantities of seeds species used in the mixture incorporated into soil material.

Legume/Grass	Quantity (kg ha ⁻¹)
<i>Zea mays</i> (L.)	24
<i>Canavalia ensiformes</i> (L)	16
<i>Helianthus annuus</i> (L.)	8
<i>Ricinus communis</i> (L)	5
<i>Calopogonium mucunoides</i>	4
<i>Pennisetum glaucum</i> (L.)	2
<i>Crotalaria juncea</i> (L.)	10
<i>Crotalaria spectabilis</i>	10
<i>Mucuna pruriens</i> (L.)	16
<i>Cajanus cajan</i> (L)	16
<i>Lablab purpureus</i> (L.)	12
<i>Sorghum bicolor</i> (L.)	3
<i>Phaseolus vulgaris</i> (L.)	12

Vegetables like lettuce, chive, coriander, rocket, pepper and tomato were grown in the ‘smart beds’. Some fruit trees like guava and Barbados cherry were also planted in the area, and some halophyte species like salt grass (*Atriplex nummularia*) and seedlings for reforestation. Aiming the animal husbandry, and considering the higher water demand and the high tolerance of grasses to salinity, it was built a breeding for planting elephantgrass (Figure 2B) irrigated with a furrow irrigation system to avoid the saline waste to be adduced into the river.

The surplus production was marketed in the community and in the market-place for family farming in neighbor communities. The resources from the sale of surplus production are used to pay for the production costs.

In order to prove the technical and environmental feasibility of using saline waste water, soil and water analysis and characterization were performed in the area according to the methodology proposed by Embrapa (1997). Besides, vegetable yield (vegetables,

P0074

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ “Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil”. EFITA-WCCA-CIGR Conference “Sustainable Agriculture through ICT Innovation”, Turin, Italy, 24-27 June 2013.

Sustainable Agriculture through ICT innovation

legumes and grasses mixture, and breeding) was quantified and the results compared to data in the literature on conventional grows (fresh water) (Maas, 1984, Arruda et al., 2011, Andrade et al., 2012).



Figure 2. Smart beds to grow vegetables irrigated by subsurface (A) and elephantgrass grown with furrow irrigation with waste from water desalination (B).

3. RESULTS AND DISCUSSION

The project enabled significant impacts on socio-economic, technical, scientific and environmental aspects in the communities supported by capture and treatment units for water desalination. A discussion was suggested with the families involved in the research, and they could find other alternatives for the reuse of saline waste, such as raising tilapia, building evaporation pan to produce minerals, and growing sunflower for oil production and ornamentation (planted in jars). During the experiment, it was found that sunflowers had higher tolerance to salinity from the water of saline waste in relation to the other species of the mixture.

From this finding, those involved in the participatory research discussed the potential and economic importance of sunflower, and further investigation was set aiming to reuse saline waste. In Table 2, it was showed the mean yield of species grown with saline waste and its relation with the conventional systems (fresh water). In relation to the level of crude protein of elephant grass, it was found an increase with the use of saline waste, but with losses in the percentage of dry matter.

It was found that there were losses in the yield of all species grown with water of saline waste, between 25.30 to 11.04% in relation to the growths using fresh water for irrigation (Table 2). However, once the vegetables yield losses are due to the effects of salinity of the saline waste, they are certainly compensatory for the gains made by the

P0074

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ “Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil”. EFITA-WCCA-CIGR Conference “Sustainable Agriculture through ICT Innovation”, Turin, Italy, 24-27 June 2013.

Sustainable Agriculture through ICT innovation

option of disposing the desalination waste appropriately, avoiding environmental impacts for disposing it in soil and water course.

Studies on the tolerance of various species to salinity have shown that, through adequate water management and cultivation practices, crops can be produced commercially with saline water (Caruso and Villari, 2004, Savvas et al., 2007; Al-Karaki et al., 2009). Thus, in soilless, it is expected that crops, provide a sustainable use for the reject brine and also, a guaranteed food production in the communities where the desalination plants were implanted.

Table 2. Mean yield of main species grown with saline waste and quality of elephantgrass (*Pennisetum purpureum* L.) forage in relation to crops grown with water of good quality (conventional system)

Species	Saline waste	Fresh water	Relative loss (%)
Lettuce (fresh biomass)	72 g per plant ⁻¹	98 g plant ⁻¹	26.53
Tomato (number of fruits)	27 fruits per plant ⁻¹	36 fruits per plant ⁻¹	25.00
Pepper (fruit weight)	43.1 g per fruit ⁻¹	57.7 g per fruit ⁻¹	25.30
Sunflower (dry biomass)	7.8 ton per ha ⁻¹	6.39 ton per ha ⁻¹	18.00
<i>Pennisetum purpureum</i> :			
- Percentage of dry matter	80.5%	90.5%	11.04
- Level of crude protein	9.5%	4%	-
Salt grass	8.31 ton per ha ⁻¹	-	-
Sabiá	10.4 g per plant ⁻¹	4.9 g per plant ⁻¹	52,74
Jurema preta	6.76 g per plant ⁻¹	4.3 g per plant ⁻¹	35,93

Still referring to Table 2, it can be seen only 11.04% reduction in the percentage of dry elephant grass and an increase in protein content when grown with saline water. According Viegas and Camarão (1994), the crude protein content should be above 7% for forage plant maintain the smooth functioning of physiological processes and also the minimum for that ruminants have a proper rumen fermentation. For herb salt (*Atriplex nummularia*), there is high dry matter production in saline environments, probably because it is a halophyte species that besides high tolerance to soil salinity, water accumulates large amounts of salts in their tissues, recovering the saline soils.

In response of Caatinga species under irrigation with water brine (Sabiá - *Mimosa caesalpinifolia* Benth e Jurema preta - *Mimosa tenuiflora*), observed in young plants reduction in dry weight of plants in relation to irrigation with good quality water in the

P0074

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ "Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil". EFITA-WCCA-CIGR Conference "Sustainable Agriculture through ICT Innovation", Turin, Italy, 24-27 June 2013.

Sustainable Agriculture through ICT innovation

order of 35.93 and 52.74% for the species jurema black and thrush. The production of seedlings of forest essences is a viable alternative for the use of the water of I reject of the desalinization, being one in the ways of destiny of Possible That residue.

4. CONCLUSION

It is concluded that there were yield losses in all species grown with water of saline waste – from 25.30% to 26.7% in relation to the conventional irrigation systems (fresh water); but there was biomass. However, the vegetable growth was successful in the community, because it could change the resident's mind about the disposal and reuse of the saline waste.

5. REFERENCES

- Al-Karaki, G., Al-Ajmi, A., Othman, Y., 2009. Response of Soilless Grown Bell Pepper Cultivars to Salinity. *Acta Horticulturae* 807(2):227- 232.
- Andrade, L.O., Cheyi, H R., Dias, N.S., Nobre, R.G., Nascimento, E.C.S., 2012. Qualidade de flores de girassóis ornamentais irrigados com águas residuária e de abastecimento. *Idesia* 30(1):19-27.
- Arruda, C.E.M., Dias, N.S., Blanco, F.F., Sousa Neto, O.N., Ferreira Neto, M., 2011. Bell pepper cultivation with brine from brackish water desalination. *Revista Caatinga* 24(2):197-201.
- Caruso, G., Villari, G., 2004. Effect of EC-level and plant shading on the NFT-grown "Friariello Pepper". *Acta Horticulturae* 659(2):576-585.
- Dias, N.S., Lima, F.A., Silva, C.R., Sousa Neto, O.N., Cheyi, H.R., 2011. Use of reject brine from desalination on different development stages of hydroponic lettuce. *Revista Caatinga* 24(1):76-81.
- Embrapa. Manual de métodos de análise de solo. 1. ed. Rio de Janeiro: EMBRAPA-CNPS, 1997, 212p.
- Fritzmman, C. et al., 2007. State-of-the-art of reverse osmosis desalination. *Desalination* 216(1):1-76.
- Maas, E.V., 1984. *Crop Tolerance*. California Agriculture 36 (1), 20 –21.
- Mohamed, A.M.O., Maraqa, M., Al Handhaly, J, 2005. Impact of land disposal of reject brine from desalination plants on soil and groundwater. *Desalination* 182(1):411-433.

P0074

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ "Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil". EFITA-WCCA-CIGR Conference "Sustainable Agriculture through ICT Innovation", Turin, Italy, 24-27 June 2013.



Sustainable Agriculture through ICT innovation

Porto, E.R., Amorim, M.C.C., Silva Junior, L.G. de A., 2011. Uso do rejeito da dessalinização de água salobra para irrigação da erva-sal (*Atriplex nummularia*). *Revista Brasileira de Engenharia Agrícola e Ambiental* 5(1):111-114.

P0074

F.I. de Souza, N. da S. Dias, O. N. de Sousa Neto, A. C. M. de Souza¹, R. B. de Lira and F. S. de Souza Junior¹ “Vegetable Growth with Wastewater from the Salt Water Treatment Station in the Community Bom Jesus, RN Brazil”. EFITA-WCCA-CIGR Conference “Sustainable Agriculture through ICT Innovation”, Turin, Italy, 24-27 June 2013.